

DPNU-94-04
January 1994

Critical Behavior of Black Hole Formation in a Scalar Wave Collapse

Yoshimi OSHIRO,¹ Kouji NAKAMURA²
and Akira TOMIMATSU³

*Department of Physics, Nagoya University
Chikusa-ku, Nagoya 464-01, Japan*

We present a self-similar model of spherically symmetric collapse of a massless scalar field with a parameter p . The black hole formation is explicitly shown to occur only in the strong-field implosion of $p > 1$. The field evolution in the critical limit $p \rightarrow 1$ is compared with numerical results found by Choptuik.

Gravitational collapse is one of interesting phenomena in general relativity. In recent years, a large number of studies have been made on the dynamical behavior of a massless scalar field, in the context of spherically symmetric gravitational collapse. It has been shown that for a weak-field implosion scalar waves bounce and disperse to infinity, while a strong-field implosion leads to black hole formation [1]. Furthermore, in numerical calculations, Choptuik [2] found an important behavior of a family of solutions which contain the parameter p characterizing the strength of the scalar field: For black hole formation there exists the critical limit $p \rightarrow p^*$ to give the gravitational mass satisfying a power law

¹e-mail: yoshiro@allegro.phys.nagoya-u.ac.jp

²e-mail: kouchan@allegro.phys.nagoya-u.ac.jp

³e-mail: c42615a@nucc.cc.nagoya-u.ac.jp

$M_{BH} \propto |p - p^*|^\gamma$ with a universal critical exponent $\gamma \simeq 0.37$. Such a critical behavior may be a general property of gravitational collapse, since imploding axisymmetric gravitational waves also show the similar behavior [3].

In this letter our purpose is to present an analytical model of the spherically symmetric collapse of scalar waves. We solve the spherically symmetric Einstein equations coupled to a massless scalar field ψ to find the self-similar solution which has a critical parameter p . This model allows us to see explicitly the black hole formation and the scalar field dispersion. Our main concern is the field evolution in the critical limit $p \rightarrow 1$, which should be compared with the numerical results.

Let us impose a self-similarity on the massless scalar field such that $\psi(u, v) = \psi(u/v)$, where u and v are retarded and advanced times respectively. The line element in the spherical symmetric double-null coordinates is given by

$$ds^2 = -h du dv + r^2 d\Omega^2, \quad (1)$$

and we assume the metric to be $h(u, v) = h(u/v)$ and $r(u, v) = v f(u/v)$. For the energy-momentum tensor of the massless scalar field defined by

$$T_{\mu\nu} = \frac{1}{4\pi} (\partial_\mu \psi \partial_\nu \psi - \frac{1}{2} g_{\mu\nu} \partial^\alpha \psi \partial_\alpha \psi), \quad (2)$$

the Einstein equations $G_{\mu\nu} = 8\pi T_{\mu\nu}$ give the unique solution as follows,

$$h = 1, \quad (3)$$

$$r^2 = \frac{1}{4} \{ (1 - p^2) v^2 - 2vu + u^2 \}, \quad (4)$$

$$\psi = \pm \frac{1}{2} \log \frac{(1 - p)v - u}{(1 + p)v - u}, \quad (5)$$

where the constant p can be chosen to be positive without loss of generality. In this letter we use units such that $G = c = 1$.

The space-time is asymptotically flat at the past null infinity $u \rightarrow -\infty$, where $r \simeq -u/2$. The scalar waves implode from the past null infinity with $\psi \simeq \pm pv/u \simeq \mp pv/r$. The parameter p turns out to give the amplitude of the imploding scalar waves. This solution is useful to analyze the scalar wave collapse, if a pathological structure due to naked singularities is removed from the space-time. Because the Ricci scalar R has the form

$$R = \frac{p^2 uv}{2r^4}, \quad (6)$$

the curvature singularity is located at the origin $r = 0$. Note that the locus of $r = 0$ is given by $u = (1 - p)v$ and $u = (1 + p)v$ in the regions $v > 0$ and $v < 0$, respectively. The space-time structure is crucially dependent on the parameter p (see Fig. 1). For $p > 1$ the singularity becomes time-like in the region $v < 0$ and spacelike in the region $v > 0$, while for $0 < p < 1$ it always becomes timelike. (For $p = 1$ exactly, the curvature singularity at $r = 0$ in the region $v > 0$ becomes null (i.e., $u=0$). We do not consider this case, and we will take the critical limit $p \rightarrow 1$ by keeping the condition $p > 1$.) The important point is that the apparent horizon exists only in the strong-field implosion of $p > 1$. In fact we obtain the expansion θ for outgoing null rays given by

$$\theta = \frac{\{(1 - p^2)v - u\}}{\{(1 - p)v - u\}\{(1 + p)v + u\}} \quad (7)$$

and the apparent horizon where $\theta = 0$ is found to be $u = (1 - p^2)v$. Thus the time-like singularity becomes naked, while the space-like one is hidden by the apparent horizon (see Fig. 1). This result will be due to the gravitational mass M which can be locally defined in the spherically symmetric system as follows,

$$M \equiv \frac{r}{2}(1 - g^{\mu\nu}\partial_\mu r \partial_\nu r) = -\frac{p^2 uv}{8r}. \quad (8)$$

Near the singularity $u = (1 - p)v$ in the region $v > 0$, Eq. (8) is reduced to $M \simeq p^2(p - 1)v^2/r$. The mass turns out to be negative near the time-like

singularity ($0 < p < 1$) and positive near the space-like one ($p > 1$). Near another timelike singularity $u = (1 + p)v$ in the region $v < 0$, we obtain $M < 0$ irrespective of p .

In 2-dimensional gravity, the similar relation among the singularity, the local gravitational mass and the apparent horizon was pointed out by Hayward in the context of the cosmic censorship [4]. Note that the 2-dimensional metric can have the form $ds^2 = r^{-2}dudv$ with r given by Eq. (4) as a special case. This may suggest that the 2-dimensional dilation gravity works as a good model of the spherically symmetric 4-dimensional gravity. However, in the 4-dimensional space-time, the dynamical scattering of imploding scalar waves plays an important role for the evolution of the scalar field. The critical behaviors which will be discussed in the following are missed in the 2-dimensional model.

It is clear that the self-similar solution is not applicable to the region $v < 0$, where the naked timelike singularity $u = (1 + p)v$ appears with $M < 0$. For a realistic evolution we must avoid such a singularity to set some regular initial data. Fortunately, this is possible, since the solution requires that there is no energy flux across the $v = 0$ line (i.e. outgoing wave ψ_u equal to zero.) and the mass is equal to zero on it. We can replace the region $v < 0$ by a flat space-time, which is smoothly matched to the region $v > 0$ (see. Fig 2). For the weak-field implosion ($0 < p < 1$) shown in Fig. 2a, the region corresponding to $u > 0$ and $v > 0$ is also replaced by a flat metric, and the time-like singularity $u = (1 - p)v$ is eliminated from the space-time. (Recall that $M = 0$ on the $u = 0$ line.) This is the case that the dynamical scattering of the imploding waves can induce a complete bounce to the future null infinity $v \rightarrow \infty$. For the strong-field implosion ($p > 1$) shown in Fig. 2b, the space-like singularity $u = (1 - p)v$ is hidden by apparent horizon. This means a black hole formation by imploding scalar waves.

However, our self-similar model requires the imploding amplitude increasing like $|\psi| \sim v$ and fails to give an asymptotically flat region at the future null infinity $v \rightarrow \infty$. (The radius $r = p(p^2 - 1)^{1/2}v/2$ of the apparent horizon infinitely increases as $v \rightarrow \infty$.) Therefore, for gravitational collapse of wave packets shown in Fig. 3, the self-similarity must break down in some region denoted by $v > v_0$, where outgoing waves dominate incoming ones. Nevertheless the discussions in the following suggest that the self-similar solution is approximately valid in a strong-field region of $0 \leq v < v_0$.

We have found the crucial difference of the space-time structure between the strong-field implosion ($p > 1$) and the weak one ($0 < p < 1$). Now let us consider the critical limit $p \rightarrow 1$ for black hole formation. The numerical calculations of wave-packet collapse done by Choptuik claim that the black hole mass approaches zero in some critical limit. To discuss such a critical behavior for our self-similar model, we must estimate the gravitational mass M_h on the apparent horizon instead of the Bondi mass which can be defined only at the future null infinity $v \rightarrow \infty$. From Eq. (8) we obtain $M_h = p\sqrt{p^2 - 1}v/4$. It is interesting to compare M_h with the initial mass $M_i = p^2v/4$ at the past null infinity $u \rightarrow -\infty$. The ratio

$$\frac{M_h}{M_i} = \frac{\sqrt{p^2 - 1}}{p}. \quad (9)$$

may be interpreted as the transmission coefficient of incoming waves propagating along a fixed v . For the strong-field limit $p \gg 1$ the amount of dynamical scattering which generates outgoing waves remains negligibly small before incoming waves reach the apparent horizon. For the critical limit $p \rightarrow 1$, however, almost all the incoming waves are scattered away, and a small portion of the generated outgoing waves can fall into the apparent horizon. (Recall that the surface $u = (1 - p^2)v$ becomes nearly null.) Thus the black hole mass M_h estimated on

the apparent horizon vanishes in the limit $p \rightarrow 1$, which gives the power law

$$M_h \simeq \left(\frac{p-1}{8}\right)^{\frac{1}{2}} v. \quad (10)$$

This is slightly different from the power law with the critical exponent $\gamma^* \simeq 0.37$, which was numerically evaluated by Choptuik at the future null infinity $v \rightarrow \infty$. We wish to emphasize that Eq. (10) exhibits a mass evolution in some dynamical stage of gravitational collapse instead of the asymptotic stage $v \rightarrow \infty$. If the self-similarity approximately holds in the stage $0 < v < v_0$ even for the wave-packet collapse shown in Fig. 3, the final mass will be given by Eq. (10) with v_0 in place of v . Then, to obtain the critical exponent $\gamma^* \simeq 0.37$, the advanced time v_0 should increase like $v_0 \propto |p-1|^{-0.13}$ in the limit $p \rightarrow 1$. For the near-critical evolution the self-similar stage may continue for a long time. Though this problem of the critical exponent remains unclear within the framework of the present analysis, we can point out another critical behavior in relation to the numerical result.

Now let us introduce the time coordinate t defined by

$$\log(t/r) = - \int \left[\frac{\sqrt{p^2 X^2 + 4} - p^2 X}{2 - X\sqrt{p^2 X^2 + 4} + p^2 X^2} \right] dX, \quad (11)$$

where $X \equiv v/r$. The line element can be rewritten into the diagonal form

$$ds^2 = -\alpha(X)dt^2 + \beta(X)dr^2 + r^2 d\Omega^2, \quad (12)$$

where the metric coefficients $\alpha(X)$ and $\beta(X)$ are

$$\alpha(X) = \left(1 - \frac{p^2 X}{\sqrt{p^2 X^2 + 4}}\right) \left[\frac{2 - X\sqrt{p^2 X^2 + 4} + p^2 X^2}{\sqrt{p^2 X^2 + 4} - p^2 X} \right]^2 \times \exp\left(2 \int \left[\frac{\sqrt{p^2 X^2 + 4} - p^2 X}{2 - X\sqrt{p^2 X^2 + 4} + p^2 X^2} \right] dX\right), \quad (13)$$

$$\beta(X) = \frac{4}{\sqrt{p^2 X^2 + 4}(\sqrt{p^2 X^2 + 4} - p^2 X)}, \quad (14)$$

The scalar field ψ also depends on X only,

$$\psi = \pm \log \frac{1}{4} (\sqrt{p^2 X^2 + 4} + pX). \quad (15)$$

We consider the field evolution measured by the coordinates t and r , which are related to the variable X through Eq. (11). The variable X satisfies $X^2 < X_h^2 \equiv 4/p^2(p^2 - 1)$ outside of the apparent horizon. Then, in the critical limit $p \rightarrow 1$, there exists the strong-field region $1 \ll X^2 \ll X_h^2$ in a neighborhood of $r = 0$, where Eq. (11) gives the logarithmic relation

$$\frac{1}{(p-1)X^2} = \log t - \log r. \quad (16)$$

Such a logarithmic evolution was also observed in the numerical calculation of wave-packet collapse with a scaling relation. Note that the time “ t ” proportional to the central proper time which defined by $T \equiv \int \sqrt{|g_{tt}(t, 0)|} dt \simeq \sqrt{|p-1|} t$. For the self-similar solution the scaling relation $\psi(\log r - \Delta, \log t - \Delta) = \psi(\log r, \log t)$ clearly hold with a continuous parameter Δ . For the wave-packet collapse, however, the scaling parameter Δ was found to be restricted to some discrete values $n\Delta^*$ ($\Delta^* \simeq 3.4$ and $n = 1, 2, 3, \dots$). This difference of the scaling will not be crucial, if we consider the region where both $\log r$ and $\log t$ are much larger than Δ^* . The apparent horizon can be arbitrarily close to $r = 0$ in the limit $p \rightarrow 1$. Then the relation $\log r \gg \Delta^*$ is satisfied in the strong-field region outside the apparent horizon near $r = 0$, where the spacing of the discrete values $n\Delta^*$ becomes unimportant to recover approximately the self-similarity.

In summary, our self-similar model can give a simple picture of black hole formation due to imploding scalar waves with the transmission coefficient (9). We obtain the critical behaviors which exhibit the mass evolution (10) on the apparent horizon and the logarithmic evolution (16) in a neighborhood of $r = 0$. The scaling relation means that the self-similarity may appear even in wave-packet collapse with the parameter p sufficiently close to the critical value and can represent a dynamical growth of the apparent horizon, though the critical

exponent γ^* and the scaling constant Δ^* are missed. It is necessary to check the self-similar critical behaviors in numerical calculations of wave-packet collapse.

Acknowledgments

This work is partially supported by the Grant-in-Aid for Scientific Research from the Ministry of Education, Science and Culture (04640268).

References

- [1] D.Chirstodoulou Commun. Math. Phys. **109**, (1987), 613; Commun. Math. Phys. **106**, (1986), 587; Commun. Math. Phys. **105**, (1986), 337:
D.S.Goldwirth and T.Piran, Phys. Rev. **D 36**, (1987), 3575.
- [2] M.W.Choptuik, Phys. Rev. Lett. **70**, (1993), 9.
- [3] A.M.Abrahams, Phys. Rev. Lett. **70**, (1993), 2980.
- [4] S.A.Hayward,Class. Quantum Grav. **10** , (1993), 985.

Figures




Fig. 1.

Penrose diagram for the self-similar solutions covering the whole space-time. The scalar waves implode from the past null infinity $\mathcal{J}^-$ with the amplitude denoted by the parameter p : (a) For the weak-field implosion $0 < p < 1$ the time-like singularity appears in the negative mass regions of $uv > 0$. (b) For the strong-field implosion $p > 1$ the apparent horizon (A.H.) is formed to surround the space-like singularity in the positive mass region. This space-time has no future null infinity $\mathcal{J}^+$ at $v \rightarrow \infty$.




Fig. 2.

Penrose diagram for the self-similar solution, in which the negative mass region are replaced by the flat space-time with $M = 0$.

(a) For $0 < p < 1$ all the imploding waves bounce and disperse to the future null infinity $\mathcal{J}^+$. (b) For $p > 1$ we obtain a model of black hole formation due to the scalar wave collapse, though the mass infinitely increases as $v \rightarrow \infty$.




Fig. 3

Penrose diagram for black hole formation due to wave-packet collapse. The region I ($v < 0$) is the flat space-time, and in the region III ($v > v_0$) outgoing waves dominate incoming ones. The self-similarity will remain valid in the region II ($0 < v < v_0$).

This figure "fig1-1.png" is available in "png" format from:

<http://arXiv.org/ps/gr-qc/9402017v3>

This figure "fig1-2.png" is available in "png" format from:

<http://arXiv.org/ps/gr-qc/9402017v3>

This figure "fig1-3.png" is available in "png" format from:

<http://arXiv.org/ps/gr-qc/9402017v3>